

兔抗 TCN2 多克隆抗体

中文名称：兔抗 TCN2 多克隆抗体

英文名称：Anti-TCN2 rabbit polyclonal antibody

别名：II; TC; TC2; TC-2; TCII; TC II; D22S676; D22S750

相关类别：一抗

抗原：TCN2

储存：冷冻（-20℃）避光

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of the vitamin B12-binding protein family. This family of proteins, alternatively referred to as R binders, is expressed in various tissues and secretions. This plasma protein binds cobalamin and mediates the transport of cobalamin into cells. This protein and other mammalian cobalamin-binding proteins, such as transcobalamin I and gastric intrinsic factor, may have evolved by duplication of a common ancestral gene. Alternative splicing results in multiple transcript variants.
Applications:	WB, IHC
Name of antibody:	TCN2
Immunogen:	Fusion protein of human TCN2
Full name:	transcobalamin II

Synonyms :	II; TC; TC2; TC-2; TCII; TC II; D22S676; D22S750
SwissProt:	P20062
IHC positive control:	Human liver cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	48 kDa
WB Positive control:	Human bladder carcinoma tissue
WB Recommended dilution:	500-2000



