

兔抗 TIA1 多克隆抗体

中文名称: 兔抗 TIA1 多克隆抗体

英文名称: Anti-TIA1 rabbit polyclonal antibody

名: 别 WDM; TIA-1

宿 主: Rabbit

抗 原: TIA1

储 存: 冷冻(-20℃)

相关类别: 一抗

标记物: Unconjugate

克隆类型: rabbit polyclonal

反应种属: Human, Mouse

技术规格

Background:	The product encoded by this gene is a member of a RNA-binding protein family and possesses nucleo lytic activity against cytotoxic lymphocyte (CTL) tar get cells. It has been suggested that this protein may be involved in the induction of apoptosis as it preferentially recognizes poly(A) homopolymers and induces DNA fragmentation in CTL targets. The maj or granule-associated species is a 15-kDa protein t hat is thought to be derived from the carboxyl ter minus of the 40-kDa product by proteolytic proces sing. Alternative splicing resulting in different isofor ms of this gene product has been described in the literature.
Applications:	ELISA, WB, IHC
Name of antibody:	TIA1



www.shjning.com

Immunogen:	Full length fusion protein
Full name:	TIA1 cytotoxic granule-associated RNA binding prot ein
Synonyms:	WDM; TIA-1
SwissProt:	P31483
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer and human tonsil
IHC Recommend dilution:	40-250
WB Predicted band size:	43 kDa
WB Positive control:	HEPG2 and NIH/3T3 cell lysates
WB Recommended dilution:	1000-5000







