

兔抗 TK1(Ab-13) 多克隆抗体

中文名称：兔抗 TK1(Ab-13) 多克隆抗体

英文名称：Anti-TK1(Ab-13) rabbit polyclonal antibody

别 名：TK2

相关类别：一抗

储 存：冷冻（-20℃）避光

宿 主：Rabbit

抗 原：TK1(Ab-13)

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Thymidine kinases play a critical role in generating the DNA synthetic precursor deoxythymidine triphosphate (dTTP) by catalyzing the phosphotransfer of phosphate from ATP to deoxythymidine (dT) and thymidine (T) in the cell. TK1 expression and activity is regulated in a cell cycle-dependent manner, accumulating during G1-phase to peak levels in S-phase before being degraded prior to cell division. Stability, but not activity, may be regulated via phosphorylation of TK1 at Ser13 by Cdc2 and/or Cdk2, but the precise mode of regulation remains elusive. These observations indicate that TK1 might be a useful marker of cell proliferation; however, recent studies have shown that TK1 plays a more significant role in the DNA damage response.
Applications:	WB, IHC, IF
Name of antibody:	TK1(Ab-13)

Immunogen:	Synthesized non-phosphopeptide derived from human TK around the phosphorylation site of serine 13 (P-G-S(p)-P-S).
Full name:	thymidine kinase 1, soluble
Synonyms :	TK2
SwissProt:	P04183
IHC positive control:	Human breast carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	25 kDa
WB Positive control:	COLO205 cells lysate
WB Recommended dilution:	500-3000
IF positive control:	HepG2 cells
IF Recommend dilution:	100-500



