

兔抗 PPIF 多克隆抗体

中文名称: 兔抗 PPIF 多克隆抗体

英文名称: Anti-PPIF rabbit polyclonal antibody

抗原: PPIF

别 名: CYP3; CypD; CyP-M; Cyp-D

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIas es catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is part of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of apoptotic and necrotic cell de ath.
Applications:	ELISA, WB, IHC
Name of antibody:	PPIF
Immunogen:	Fusion protein of human PPIF
Full name:	peptidylprolyl isomerase F



www.shjning.com

Synonyms:	CYP3; CypD; CyP-M; Cyp-D
SwissProt:	P30405
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human breast cancer and Human liver cancer
IHC Recommend dilution:	100-300
WB Predicted band size:	22 kDa
WB Positive control:	Hela cells
WB Recommended dilution:	500-2000





