

## 兔抗 TCF4/12 多克隆抗体

中文名称：兔抗 TCF4/12 多克隆抗体

英文名称：Anti-TCF4/12 rabbit polyclonal antibody

别名：E2-2; ITF2; PTHS; SEF2; ITF-2; SEF-2; TCF-4; SEF2-1; SEF2-1A; SEF2-1B; SEF2-1D; bHLHb19/HEB; CRS3; HTF4; bHLHb20; HsT17266

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：TCF4/12

反应种属：Human Mouse Rat

标记物：Unconjugate

克隆类型：Unconjugate

### 技术规格

<b>Background:</b>	This gene encodes transcription factor 4, a basic helix-loop-helix transcription factor. The encoded protein recognizes an E-box ('E-box') binding site ('CANNTG') - a motif first identified in immunoglobulin enhancers. This gene is broadly expressed, and may play an important role in nervous system development. Defects in this gene are a cause of Pitt-Hopkins syndrome. Multiple alternatively spliced transcript variants that encode different proteins have been described.
<b>Applications:</b>	WB
<b>Name of antibody:</b>	TCF4/12
<b>Immunogen:</b>	Synthesized peptide derived from internal of human TCF4/12.

<b>Full name:</b>	transcription factor 4/12
<b>Synonyms :</b>	E2-2; ITF2; PTHS; SEF2; ITF-2; SEF-2; TCF-4; SEF2-1; SEF2-1A; SEF2-1B; SEF2-1D; bHLHb19/HEB; CRS3; H TF4; bHLHb20; HsT17266
<b>SwissProt:</b>	P15884/Q99081
<b>WB Predicted band size:</b>	71 kDa
<b>WB Positive control:</b>	Huvec cells lysate
<b>WB Recommended dilution:</b>	500-3000

