

## ERG 抗原（重组蛋白）

中文名称：ERG 抗原（重组蛋白）

英文名称：ERG Antigen (Recombinant Protein)

储 存：冷冻（-20℃）

相关类别：抗原

概述

Fusion protein corresponding to a region derived from N terminal 150 amino acids of human ERG

技术规格

<b>Full name:</b>	v-ets erythroblastosis virus E26 oncogene homolog (avian)
<b>Synonyms:</b>	p55; erg-3
<b>Swissprot:</b>	P11308
<b>Gene Accession:</b>	BC040168
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	This gene encodes a member of the erythroblast transformation-specific (ETS) family of transcription factors. All members of this family are key regulators of embryonic development, cell proliferation, differentiation, angiogenesis, inflammation, and apoptosis. The protein encoded by this gene is mainly expressed in the nucleus. It contains an ETS DNA-binding domain and a PNT (pointed) domain which is implicated in the self-association of chimeric oncoproteins. This protein is required for platelet adhesion to the subendothelium, inducing vascular cell remodeling. It also regulates hematopoiesis, and the differentiation and maturation of megakaryocytic cells. This gene is involved in chromosomal translocations, resulting in different fusion gene products, such as TMPSSR2-ERG and NDRG1-ERG in prostate cancer, EWS-ERG in Ewing's sarcoma and FUS-ERG in acute myeloid leukemia. Multiple altern

actively spliced transcript variants encoding different isoforms have been identified.