

CSTA 抗原（重组蛋白）

中文名称：CSTA 抗原（重组蛋白）

英文名称：CSTA Antigen (Recombinant Protein)

别名：AREI; STF1; STFA

储存：冷冻（-20℃）

相关类别：抗原

概述

Full length fusion protein

技术规格

Full name:	cystatin A (stefin A)
Synonyms:	AREI; STF1; STFA
Swissprot:	P01040
Gene Accession:	BC010379
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and dia

gnostic tools for cancer.