

## CXCL14 抗原（重组蛋白）

中文名称：CXCL14 抗原（重组蛋白）

英文名称：CXCL14 Antigen (Recombinant Protein)

储存：冷冻（-20℃）

相关类别：抗原

概述

Full length fusion protein

技术规格

<b>Full name:</b>	chemokine (C-X-C motif) ligand 14
<b>Synonyms:</b>	KEC; KS1; BMAC; BRAK; NJAC; MIP2G; MIP-2g; SCYB14
<b>Swissprot:</b>	O95715
<b>Gene Accession:</b>	BC003513
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	This gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXCL (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation.