

DCPS 抗原(重组蛋白)

中文名称: DCPS 抗原 (重组蛋白)

英文名称: DCPS Antigen (Recombinant Protein)

别名: decapping enzyme, scavenger; ARS; DCS1; HSL1; HINT5; HINT-5; HSPC015

储存: 冷冻(-20℃)

相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 2-337 amino acids of human DCPS

技术规格

Full name:	decapping enzyme, scavenger
Synonyms:	ARS; DCS1; HSL1; HINT5; HINT-5; HSPC015
Swissprot:	Q96C86
Gene Accession:	BC014532
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	Decapping scavenger enzyme that catalyzes the cleavage of a residual cap structure following the degradation of mRNAs by the 3'->5' exos ome-mediated mRNA decay pathway. Hydrolyzes cap analog structure s like 7-methylguanosine nucleoside triphosphate (m7GpppG) with up to 10 nucleotide substrates (small capped oligoribonucleotides) and sp ecifically releases 5'-phosphorylated RNA fragments and 7-methylguan osine monophosphate (m7GMP). Cleaves cap analog structures like tri- methyl guanosine nucleoside triphosphate (m3(2,2,7)GpppG) with very poor efficiency. Does not hydrolyze unmethylated cap analog (GpppG) and shows no decapping activity on intact m7GpppG-capped mRNA



www.shjning.com

molecules longer than 25 nucleotides. Does not hydrolyze 7-methylgu anosine diphosphate (m7GDP) to m7GMP (PubMed:22985415). May als o play a role in the 5'->3 mRNA decay pathway; m7GDP, the downstr eam product released by the 5'->3' mRNA mediated decapping activit y, may be also converted by DCPS to m7GMP (PubMed:14523240). Bi nds to m7GpppG and strongly to m7GDP. Plays a role in first intron s plicing of pre-mRNAs. Inhibits activation-induced cell death.