

兔抗 TSC22D1 多克隆抗体

中文名称: 兔抗 TSC22D1 多克隆抗体

英文名称: Anti-TSC22D1 rabbit polyclonal antibody

别 名: Ptg-2; TSC22; TGFB1I4

相关类别: 一抗

储 存: 冷冻 (-20℃) 避光

宿 主: Rabbit

抗原: TSC22D1

反应种属: Human Mouse Rat

标记物: Unconjugate

克隆类型: Unconjugate


技术规格

ļ	,
Background:	This gene encodes a member of the TSC22 domain family of leucine zipper transcription factors. The encoded protein is stimulated by transforming growth factor beta, and regulates the transcription of multiple genes including C-type natriuretic peptide. The encoded protein may play a critical role in tumor suppression through the induction of cancer cell apoptosis, and a single nucleotide polymorphism in the promoter of this gene has been associated with diabetic nephropathy. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
Applications:	WB, IHC, IF
• •	TSC22D1
Name of antibody:	
Immunogen:	Synthesized peptide derived from internal of human TSC2



www.shjning.com

	2D1.
Full name:	TSC22 domain family, member 1
Synonyms:	Ptg-2; TSC22; TGFB1I4
SwissProt:	Q15714
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	110 kDa
WB Positive control:	Mouse liver tissue and Rat liver tissue lysates
WB Recommended dilution:	500-3000
IF positive control:	HepG2 cells
IF Recommend dilution:	100-500





www.shjning.com

