

## 兔抗 UBE2D2 多克隆抗体

- 中文名称：兔抗 UBE2D2 多克隆抗体
- 英文名称：Anti-UBE2D2 rabbit polyclonal antibody
- 别名：UBC4; PUBC1; UBCH4; UBC4/5; UBCH5B; E2(17)KB2
- 相关类别：一抗
- 储存：冷冻（-20℃）避光
- 宿主：Rabbit
- 抗原：UBE2D2
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：Unconjugate

### 技术规格

**Background:**

Regulated degradation of misfolded, damaged or short-lived proteins in eukaryotes occurs via the ubiquitin (Ub)-proteasome system (UPS). An integral part of the UPS system is the ubiquitination of target proteins and covalent linkage of Ub-containing proteins to form polymeric chains, marking them as targets for 26S proteasome-mediated degradation. Ubiquitination of proteins is mediated by a cascade of enzymes which includes E1 (ubiquitin activating), E2 (ubiquitin conjugating), and E3 (ubiquitin ligases) enzymes. This gene encodes a member of the E2 enzyme family. Substrates of this enzyme include the tumor suppressor protein p53 and peroxisomal biogenesis factor 5 (PEX5). Alternative splicing results in multiple

	e transcript variants of this gene.
<b>Applications:</b>	WB
<b>Name of antibody:</b>	UBE2D2
<b>Immunogen:</b>	Synthesized peptide derived from C-terminal of human UBE2D2.
<b>Full name:</b>	ubiquitin-conjugating enzyme E2D 2
<b>Synonyms :</b>	UBC4; PUBC1; UBCH4; UBC4/5; UBCH5B; E2(17)KB2
<b>SwissProt:</b>	P62837
<b>WB Predicted band size:</b>	17 kDa
<b>WB Positive control:</b>	Hela cells, A549 cells and K562 cells lysates
<b>WB Recommended dilution:</b>	500-3000

