

## 兔抗 UBE2D3 多克隆抗体

中文名称：兔抗 UBE2D3 多克隆抗体

英文名称：Anti-UBE2D3 rabbit polyclonal antibody

别名：UBC4/5; UBCH5C; E2(17)KB3

储存：冷冻（-20℃）

宿主：Rabbit

抗原：UBE2D3

反应种属：Human, mouse, Rat

相关类别：一抗

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme functions in the ubiquitination of the tumor-suppressor protein p53, which is induced by an E3 ubiquitin-protein ligase. Multiple spliced transcript variants have been found for this gene, but the full-length nature of some variants has not been determined.
<b>Applications:</b>	ELISA, WB, IHC

<b>Name of antibody:</b>	UBE2D3
<b>Immunogen:</b>	Full length fusion protein
<b>Full name:</b>	ubiquitin conjugating enzyme E2D 3
<b>Synonyms:</b>	UBC4/5; UBCH5C; E2(17)KB3
<b>SwissProt:</b>	P61077
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human colorectal cancer and human tonsil
<b>IHC Recommend dilution:</b>	25-100
<b>WB Predicted band size:</b>	17 kDa
<b>WB Positive control:</b>	NIH/3T3 cell lysate
<b>WB Recommended dilution:</b>	200-1000



