

兔抗 VTI1A 多克隆抗体

中文名称：兔抗 VTI1A 多克隆抗体

英文名称：Anti-VTI1A rabbit polyclonal antibody

别名：MMDS3; MVti1; VTI1RP2; Vti1-rp2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：VTI1A

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a member of the family of soluble N-ethylmaleimide-sensitive fusion protein-attachment protein receptors (SNAREs) that function in intracellular trafficking. This family member is involved in vesicular transport between endosomes and the trans-Golgi network. It is a vesicle-associated SNARE (v-SNARE) that interacts with target membrane SNAREs (t-SNAREs). Polymorphisms in this gene have been associated with binocular function, and also with susceptibility to colorectal and lung cancers. A recurrent rearrangement has been found between this gene and the transcription factor 7-like 2 (TCF7L2) gene in colorectal cancers. Alternative splicing results in multiple transcript variants.
Applications:	ELISA, WB, IHC

Name of antibody:	VTI1A
Immunogen:	Full length fusion protein
Full name:	vesicle transport through interaction with t-SNAREs 1A
Synonyms:	MMDS3; MVti1; VTI1RP2; Vti1-rp2
SwissProt:	Q96AJ9
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	25 kDa
WB Positive control:	231,Jurkat,Hela and HEPG2 cell lysates
WB Recommended dilution:	200-1000



