

兔抗 VWA5A 多克隆抗体

中文名称: 兔抗 VWA5A 多克隆抗体

英文名称: Anti-VWA5A rabbit polyclonal antibody

别 名: BCSC1; BCSC-1; LOH11CR2A

相关类别: 一抗

抗原: VWA5A

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

somal region 2 gene A protein), is a 786 amino acid protein containing one VIT domain and one VWFA domain. VWA5A is expressed at low levels in various tissues, with no expression found in 80% of tumor cell lines. Like ely acting as a tumor suppressor gene, deletion of VWA5A leads to loss of heterozygosity (LOH) in breast and ovarian tumors, and may have an important role as a potential gene therapy target. Abnormal expression of VWA5A may lead to an increase in adhesion of CNE-2L2

VWA5A (von Willebrand factor A domain containing 5A), also known as BCSC-1 (breast cancer suppressor candid ate 1) or LOH11CR2A (loss of heterozygosity 11 chromo

cells associated with an increase in expression of E-cadh erin, alpha-catenin, and p53, resulting in a decrease of



	malignant activity in cells with ectopic expression of VW
	A5A.
Applications:	ELISA, IHC
Name of antibody:	VWA5A
Immunogen:	Fusion protein of human VWA5A
Full name:	von Willebrand factor A domain containing 5A
Synonyms:	BCSC1; BCSC-1; LOH11CR2A
SwissProt:	O00534
ELISA Recommended dilution:	2000-10000
IHC positive control:	Human cervical cancer and human lung cancer
IHC Recommend dilution:	100-300

