

兔抗 WASF2 多克隆抗体

中文名称：兔抗 WASF2 多克隆抗体

英文名称：Anti-WASF2 rabbit polyclonal antibody

别名：IMD2; SCAR2; WASF4; WAVE2; dJ393P12.2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：WASF2

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of the Wiskott-Aldrich syndrome protein family. The gene product is a protein that forms a multiprotein complex that links receptor kinases and actin. Binding to actin occurs through a C-terminal verprolin homology domain in all family members. The multiprotein complex serves to transduce signals that involve changes in cell shape, motility or function. The published map location (PMID:10381382) has been changed based on recent genomic sequence comparisons, which indicate that the expressed gene is located on chromosome 1, and a pseudogene may be located on chromosome X.
Applications:	ELISA, WB, IHC

Name of antibody:	WASF2
Immunogen:	Synthetic peptide of human WASF2
Full name:	WAS protein family, member 2
Synonyms :	IMD2; SCAR2; WASF4; WAVE2; dJ393P12.2
SwissProt:	Q9Y6W5
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human brain and Human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	54 kDa
WB Positive control:	Mouse lung tissue
WB Recommended dilution:	200-1000



