

小鼠抗 CYCS 单克隆抗体

中文名称: 小鼠抗 CYCS 单克隆抗体

英文名称: Anti-CYCS mouse monoclonal antibody

别 名: cytochrome c; somatic; CYC; HCS; THC4

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Mouse

抗 原: CYCS

反应种属: Human, Mouse

标 记 物: Unconjugate


克隆类型: mouse monoclonal

技术规格


Background:	This gene encodes a small heme protein that function s as a central component of the electron transport ch ain in mitochondria. The encoded protein associates wi th the inner membrane of the mitochondrion where it accepts electrons from cytochrome b and transfers the m to the cytochrome oxidase complex. This protein is also involved in initiation of apoptosis. Mutations in th is gene are associated with autosomal dominant nonsy ndromic thrombocytopenia. Numerous processed pseu dogenes of this gene are found throughout the human genome.
Applications:	WB, IHC
Name of antibody:	CYCS
Immunogen:	Fusion protein of human CYCS



Full name:	cytochrome c, somatic
Synonyms:	CYC; HCS; THC4
SwissProt:	P99999
IHC positive control:	Human tonsil and Human thyroid cancer; Human ovari an cancer and Human esophagus cancer
IHC Recommend dilution:	100-500
WB Predicted band size:	12 kDa
WB Positive control:	Hela, 231, HepG2 and Jurkat cell, Mouse kidney tissue lysates
WB Recommended dilution:	5000-20000









www.shjning.com

