

CDK2AP1 抗原(重组蛋白)

- 中文名称: CDK2AP1 抗原(重组蛋白)
- 英文名称: CDK2AP1 Antigen (Recombinant Protein)
- 别 名: DOC1; ST19; DORC1; doc-1; p12DOC-1
- 储存: 冷冻(-20℃)
- 相关类别: 抗原

概述

Full length fusion protein

技术规格

Full name:	cyclin dependent kinase 2 associated protein 1					
Synonyms:	DOC1; ST19; DORC1; doc-1; p12DOC-1					
Swissprot:	O14519					
Gene Accession:	BC034717					
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE					
Expression system:	Escherichia coli					
Tags:	His tag C-Terminus, GST tag N-Terminus					
Background:	The protein encoded by this gene is a cyclin-dependent kinas e 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a r egulatory role in DNA replication during the S-phase of the c ell cycle. This protein also forms a core subunit of the nucleo some remodeling and histone deacetylation (NURD) complex t hat epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic r					



egulation.	Alternative	splicing	results in	multiple	transcript	vari
ants enco	ding distinc	t isoform	าร.			