

F10 抗原(重组蛋白)

中文名称: F10 抗原(重组蛋白)

英文名称: F10 Antigen (Recombinant Protein)

别 名: FX; FXA

储 存: 冷冻(-20℃)

相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 235-488 amino acids of human F10

技术规格

| Full name: | Coagulation factor X |
|--------------------|--|
| Synonyms: | FX; FXA |
| Swissprot: | P00742 |
| Gene Accession: | BC046125 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | This gene encodes the vitamin K-dependent coagulation factor X of the blood coagulation cascade. This factor undergoes multiple processing steps before its preproprotein is converted to a mature two-chain form by the excision of the tripeptide RKR. Two chains of the factor are held together by 1 or more disulfide bonds; the light chain contains 2 EGF-like domains, while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. The mature factor is activated by the cleavage of the activation peptide by factor IXa (in the intrisic pathway), or by factor VIIa (in the extrinsic pathway). The activated factor then converts pr |



www.shjning.com

othrombin to thrombin in the presence of factor Va, Ca+2, and phospholipid during blood clotting. Mutations of this gene resul t in factor X deficiency, a hemorrhagic condition of variable sev erity.