

PSMB3 抗原（重组蛋白）

中文名称：PSMB3 抗原（重组蛋白）

英文名称：PSMB3 Antigen (Recombinant Protein)

别名：proteasome subunit beta 3; HC10-II

储存：冷冻（-20℃）

相关类别：抗原

概述：

Fusion protein corresponding to a region derived from 2-205 amino acids of human PSMB3

技术规格：

Full name:	proteasome subunit beta 3
Synonyms:	HC10-II
Swissprot:	P49720
Gene Accession:	BC013008
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family,

that is a 20S core beta subunit. The 26 S proteasome may be involved in trinucleotide repeat expansion, a phenomenon which is associated with many hereditary neurological diseases. Pseudogenes have been identified on chromosomes 2 and 12. Alternative splicing results in multiple transcript variants.