

RPS27A 抗原(重组蛋白)

中文名称: RPS27A 抗原(重组蛋白)

英文名称: RPS27A Antigen (Recombinant Protein)

别 名: UBC; S27A; CEP80; UBA80; HEL112; UBCEP1; UBCEP80

储 存: 冷冻 (-20℃)

相关类别: 抗原

概述:

Fusion protein corresponding to a region derived from 16-156 amino acids of human RPS27A

技术规格:

Full name:	ribosomal protein S27a
Synonyms:	UBC; S27A; CEP80; UBA80; HEL112; UBCEP1; UBCEP80
Swissprot:	P62979
Gene Accession:	BC001392
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	Ubiquitin, a highly conserved protein that has a major role in target ing cellular proteins for degradation by the 26S proteosome, is synt hesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin fused to an unrelated protein. This gene en codes a fusion protein consisting of ubiquitin at the N terminus and ribosomal protein S27a at the C terminus. When expressed in yeast, the protein is post-translationally processed, generating free ubiquitin monomer and ribosomal protein S27a. Ribosomal protein S27a is a component of the 40S subunit of the ribosome and belongs to the S27AE family of ribosomal proteins. It contains C4-type zinc fing



www.shjning.com

er domains and is located in the cytoplasm. Pseudogenes derived fr om this gene are present in the genome. As with ribosomal protein S27a, ribosomal protein L40 is also synthesized as a fusion protein with ubiquitin; similarly, ribosomal protein S30 is synthesized as a fu sion protein with the ubiquitin-like protein fubi. Multiple alternativel y spliced transcript variants that encode the same proteins have been identified.