

## SF3B3 抗原(重组蛋白)

中文名称: SF3B3 抗原(重组蛋白)

英文名称: SF3B3 Antigen (Recombinant Protein)

别 名: RSE1; SAP130; SF3b130; STAF130

储 存: 冷冻(-20℃)

相关类别: 抗原

概述:

Fusion protein corresponding to a region derived from 1008-1207 amino acids of human SF3B3

## 技术规格:

| Full name:         | splicing factor 3b subunit 3  |
|--------------------|---|
| Synonyms:          | RSE1; SAP130; SF3b130; STAF130  |
| Swissprot:         | Q15393  |
| Gene Accession:    | BC000463  |
| Purity:            | >85%, as determined by Coomassie blue stained SDS-PA<br>GE  |
| Expression system: | Escherichia coli  |
| Tags:              | His tag C-Terminus, GST tag N-Terminus  |
| Background:        | This gene encodes subunit 3 of the splicing factor 3b pr otein complex. Splicing factor 3b, together with splicing f actor 3a and a 12S RNA unit, forms the U2 small nuclear ribonucleoproteins complex (U2 snRNP). The splicing fact or 3b/3a complex binds pre-mRNA upstream of the intro n's branch site in a sequence independent manner and m ay anchor the U2 snRNP to the pre-mRNA. Splicing facto r 3b is also a component of the minor U12-type spliceos |



## www.shjning.com

ome. Subunit 3 has also been identified as a component of the STAGA (SPT3-TAF(II)31-GCN5L acetylase) transcripti on coactivator-HAT (histone acetyltransferase) complex, and the TFTC (TATA-binding-protein-free TAF(II)-containing complex). These complexes may function in chromatin m odification, transcription, splicing, and DNA repair.