

## DLAT 抗原（重组蛋白）

中文名称： DLAT 抗原（重组蛋白）

英文名称： DLAT Antigen (Recombinant Protein)

别名： dihydrolipoamide S-acetyltransferase; DLTA; PDCE2; PDC-E2

储存： 冷冻（-20℃）

相关类别： 抗原

### 概述

Fusion protein corresponding to C terminal 200 amino acids of human DLAT

### 技术规格

<b>Full name:</b>	dihydrolipoamide S-acetyltransferase
<b>Synonyms:</b>	DLTA; PDCE2; PDC-E2
<b>Swissprot:</b>	P10515
<b>Gene Accession:</b>	BC039084
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	This gene encodes component E2 of the multi-enzyme pyruvate dehydrogenase complex (PDC). PDC resides in the inner mitochondrial membrane and catalyzes the conversion of pyruvate to acetyl coenzyme A. The protein product of this gene, dihydrolipoamide acetyltransferase, accepts acetyl groups formed by the oxidative decarboxylation of pyruvate and transfers them to coenzyme A. Dihydrolipoamide acetyltransferase is the antigen for antimitochondrial antibodies. These autoantibodies are present in nearly 95% of patients with the autoimmune liver disease primary biliary cirrhosis (PBC). In PBC, activated T lymphocytes attack and destroy epithelial cells in the bile duct where this prot

ein is abnormally distributed and overexpressed. PBC eventually leads to cirrhosis and liver failure. Mutations in this gene are also a cause of pyruvate dehydrogenase E2 deficiency which causes primary lactic acidosis in infancy and early childhood.