

## RPL3 抗原（重组蛋白）

中文名称： RPL3 抗原（重组蛋白）

英文名称： RPL3 Antigen (Recombinant Protein)

别名： ribosomal protein L3; L3; ASC-1; TARBP-B

储存： 冷冻（-20℃）

相关类别： 抗原

### 概述

Fusion protein corresponding to a region derived from 204-403 amino acids of human RPL3

### 技术规格

<b>Full name:</b>	ribosomal protein L3
<b>Synonyms:</b>	L3; ASC-1; TARBP-B
<b>Swissprot:</b>	P39023
<b>Gene Accession:</b>	BC002408
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	Ribosomes, the complexes that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L3P family of ribosomal proteins and it is located in the cytoplasm. The protein can bind to the HIV-1 TAR mRNA, and it has been suggested that the protein contributes to tat-mediated transactivation. This gene is co-transcribed with several small nucleolar RNA genes, which are located in several of this gene's introns. Alternate transcriptional splice va

riants, encoding different isoforms, have been characterized. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.