

SEMA4A 抗原（重组蛋白）

中文名称：SEMA4A 抗原（重组蛋白）

英文名称：SEMA4A Antigen (Recombinant Protein)

别名：sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4A; RP35; SEMB; SEMAB;

储存：冷冻（-20℃）

相关类别：抗原

概述

Fusion protein corresponding to C terminal 201 amino acids of human SEMA4A

技术规格

Full name:	sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4A
Synonyms:	RP35; SEMB; SEMAB; CORD10
Swissprot:	Q9H3S1
Gene Accession:	BC020974
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	This gene encodes a member of the semaphorin family of soluble and transmembrane proteins. Semaphorins are involved in numerous functions, including axon guidance, morphogenesis, carcinogenesis, and immunomodulation. The encoded protein is a single-pass type I membrane protein containing an immunoglobulin-like C2-type domain, a PSI domain and a sema domain. It inhibits axonal extension by providing local signals to specify territories inaccessible for growing axons. It is an activator of T-cell-mediated immunity and suppresses vascula

	<p>r endothelial growth factor (VEGF)-mediated endothelial cell migration and proliferation in vitro and angiogenesis in vivo. Mutations in this gene are associated with retinal degenerative diseases including retinitis pigmentosa type 35 (RP35) and cone-rod dystrophy type 10 (CORD10). Multiple alternatively spliced transcript variants encoding different isoforms have been identified.</p>
--	---