

UBXN2B 抗原(重组蛋白)

UBXN2B 抗原(重组蛋白) 中文名称:

英文名称: UBXN2B Antigen (Recombinant Protein)

别 名: UBX domain protein 2B; p37

相关类别: 抗原

存: 冷冻(-20℃)

概述

Fusion protein corresponding to a region derived from 132-331 amino acids of human UBXN2B

技术规格

| Full name: | UBX domain protein 2B |
|--------------------|---|
| Synonyms: | p37 |
| Swissprot: | Q14CS0 |
| Gene Accession: | BC113645 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | UBXD2B (UBX domain-containing protein 2B), also known as N SFL1 cofactor p37 and p97 cofactor p37, is a 331 amino acid protein that contains one UBX domain and one SEP domain. U BXN2B is required for ER and Golgi biogenesis and also plays a role in their maintenance during interphase, as well as their reassembly at the end of mitosis. Through interaction with VCP, UBXN2B forms a complex that has membrane fusion activity. Adapter protein required for Golgi and endoplasmic reticulum biogenesis. Involved in Golgi and endoplasmic reticulum maint |



www.shjning.com

enance during interphase and in their reassembly at the end of mitosis. The complex formed with VCP has membrane fusion activity; membrane fusion activity requires USO1-GOLGA2 tethering and BET1L. VCPIP1 is also required, but not its deubiquitinating activity.